

DESY 97-150

August 1997

ON THE ODDERON INTERCEPT IN QCD¹

N. Armesto

*II. Institut für Theoretische Physik, Universität Hamburg,
Luruper Chaussee 149, D-22761 Hamburg, Germany*

and

M. A. Braun

*Department of High Energy Physics, University of St. Petersburg,
198904 St. Petersburg, Russia*

The odderon singularity is studied in perturbative QCD in the framework of the Bartels-Kwieciński-Praszalowicz (BKP) equation. Arguments for the odderon intercept being exactly equal to unity are given. Besides, a variational method based on a complete system of one-gluon functions is presented. For the odderon, the highest intercept calculated by this method is $1 - (N_c \alpha_s / \pi) 0.45$. Comparison to other calculations is shown.

¹Talk given by N. Armesto at the *Madrid Workshop on low x Physics* (Miraflores de la Sierra, Spain, June 18th-21st 1997).

1 Introduction

Regge theory [1] has been used during the last 30 years to describe strong interaction at high energies and low transferred momenta. The amplitude for the reaction $p_a p_b \longrightarrow p'_a p'_b$, in the limit $s \gg m^2 \simeq -t$, can be expressed as a sum over Regge trajectories $j(t) = 1 + \omega(t)$ exchanged in the t -channel:

$$A(s, t) = \sum_{j; p=\pm} \xi_{j_p(t)}^p s^{j_p(t)} g_1^p(t) g_2^p(t), \quad (1)$$

with $s = (p_a + p_b)^2$ and $t = (p_a - p'_a)^2$ the Mandelstam variables and $\xi_j^p = i - (\cos \pi j + p)/\sin \pi j$ the signature factor.

The amplitude can be decomposed as a sum over parts with definite signature, i.e. definite behaviour under the exchange $s \longrightarrow -s$:

$$A(s, t) = A^+(s, t) + A^-(s, t), \quad A^\pm(s, t) = \pm A^\pm(-s, t). \quad (2)$$

As the total cross section can be related to the amplitude via the optical theorem, $\sigma_{tot} \propto \text{Im } A(s, 0)/s$, the behaviour at high energies will be determined by those Regge trajectories exchanging vacuum quantum numbers and with highest intercept $\omega(0)+1 = j(0)$. For positive signature (contributing equally both to pp and $\bar{p}p$ scattering) the trajectory with the highest intercept is called the pomeron and its intercept has been determined [2] to be supercritical (i.e. > 1): $\omega(t) \simeq 0.08 + (0.25 \text{ GeV}^{-2}) t$. The trajectory with negative signature (contributing to the difference between pp and $\bar{p}p$) has been called the odderon [3]; phenomenological fits to soft data seem to indicate that its contribution to soft interactions at high energies is negligible, although this is still a matter of debate [4].

There have been several attempts to relate Regge theory to QCD. Within the framework of perturbative QCD, in the limit $\alpha_s \ln s \sim 1$ (leading-log approximation in s), the so-called Balitsky-Fadin-Kuraev-Lipatov [5] (BFKL) pomeron appears as a fixed cut in the j -plane with intercept $1 + \omega_{BFKL}(0) = 1 + (N_c \alpha_s / \pi) 4 \ln 2$ ($\simeq 1.5$ for $N_c = 3$ and $\alpha_s \simeq 0.2$); it corresponds to the bound state of two t -channel reggeized

gluons. Under the same approximation the odderon appears as the bound state of three gluons in a symmetric colour configuration, given by the solution of the BKP [6] equation.

Attempts to solve the odderon problem have gone in different directions. On the one hand two-dimensional conformal techniques have been applied [7, 8]. On the other hand variational methods have been used, both with conformal invariant [9, 10] and polynomial [11] trial functions. This constitutes a first step towards the unitarization [12] of QCD at high energies, which, together with next-to-leading-log corrections [13], are expected to give full consistency to the whole approach [14].

In this contribution we will treat the following aspects [11]: In Sect. 2 arguments will be presented for the odderon intercept being exactly unity. In Sect. 3 a variational method will be proposed. Finally in Sect. 4 some numerical results and comparison to other calculations will be shown.

2 Argument for intercept equal to unity

The BKP equation for three gluons with transverse momenta q_1, q_2, q_3 can be written in a Hamiltonian form with $E = 1 - j$:

$$H\psi = E\psi, \quad H = T_1 + T_2 + T_3 + U_{12} + U_{23} + U_{31}. \quad (3)$$

In units of $N_c\alpha_s/\pi$,

$$T_1 = -\omega(q_1) = \frac{\eta(q_1)}{4\pi} \int \frac{d^2q'_1}{\eta(q'_1)\eta(q_1 - q'_1)}, \quad (4)$$

with $\eta(q) = q^2 + m^2$, is the gluon Regge trajectory. In (3) the dependence on m vanishes, so the BKP equation is infrared stable [6, 15].

The U_{ik} 's are integral operators which act with measure

$$\begin{aligned} d\mu &= d^2q_1 d^2q_2 d^2q_3 \delta^{(2)}(q_1 + q_2 + q_3 - q) / [\eta(q_1)\eta(q_2)\eta(q_3)], \\ U_{12}(q_1, q_2, q_3 | q'_1, q'_2, q'_3) &= \eta(q_3) \delta^{(2)}(q_3 - q'_3) V_{12}(q_1, q_2 | q'_1, q'_2). \end{aligned} \quad (5)$$

Here the V_{ij} 's are BFKL interaction kernels for 2 gluons in a vector colour state (a factor 1/2 with respect to the vacuum channel appears):

$$V_{12}(q_1, q_2 | q'_1, q'_2) = -\frac{1}{4\pi} \left[\frac{\eta(q_1)\eta(q'_2) + \eta(q'_1)\eta(q_2)}{\eta(q_1 - q'_1)} - \eta(q_1 + q_2) \right]. \quad (6)$$

Due to the bootstrap identity [16]:

$$\int \frac{d^2 q'_1}{\eta(q'_1)\eta(q_1 + q_2 - q'_1)} V_{12}(q_1, q_2 | q'_1, q'_2) = \omega(q_1) + \omega(q_2) - \omega(q_1 + q_2), \quad (7)$$

for $q = q_1 + q_2 + q_3 = 0$, $\psi_B(q_1, q_2, q_3) = \psi_0$ is a solution with maximal symmetry which gives $E = 0$. This solution does not fulfil the gauge invariance requirement $\psi_B(q_i = 0) = 0$ and for $m \rightarrow 0$ it decouples from the physical spectrum, but still offers a lower bound for the energy (as the state with $|n| = 1$, $\nu = 0$ for the $|n| = 1$ sector in the BFKL pomeron [17]). More elaborated mathematical arguments can be found in [11].

3 Variational method

Setting $m = 0$, and redefining $\psi \rightarrow \prod_{i=1}^3 q_i^2 \psi$, we have

$$H\psi = E \prod_{i=1}^3 q_i^2 \psi, \quad H = (1/2)(H_{12} + H_{23} + H_{31}) \quad (8)$$

and, in a mixed representation ($\mathbf{C}$ is Euler's constant),

$$\begin{aligned} H_{ik}\psi &= \prod_{j=1}^3 q_j^2 (\ln q_i^2 q_k^2 + 4\mathbf{C})\psi + \prod_{j=1, j \neq i, k}^3 q_j^2 \left[q_i^2 \ln(r_{ik}^2/4) q_k^2 + (i \leftrightarrow k) \right] \psi \\ &+ 2(q_i + q_k)^2 \psi(r_i - r_k = 0). \end{aligned} \quad (9)$$

As usually, the variational approach provides us with an upper bound for the ground state energy ($\equiv$ lower bound for the intercept) and consists in finding a minimum of the functional

$$\begin{aligned} \Phi &= \int \prod d^2 q_i \psi^* H \psi, \\ \int \prod d^2 q_i \psi^* \prod_{j=1}^3 q_j^2 \psi &= 1. \end{aligned} \quad (10)$$

As trial function we choose a linear combination of one-gluon functions:

$$\psi(r_1, r_2, r_3) = \sum_{\alpha_1, \alpha_2, \alpha_3} c_{\alpha_1, \alpha_2, \alpha_3} \prod_{i=1}^3 \psi_{\alpha_i}(r_i), \quad (11)$$

with

$$\int d^2r \psi_{\alpha}^* q^2 \psi_{\alpha'} = \delta_{\alpha, \alpha'}, \quad \sum_{\alpha_1, \alpha_2, \alpha_3} |c_{\alpha_1, \alpha_2, \alpha_3}|^2 = 1 \quad (12)$$

and $c_{\alpha_1, \alpha_2, \alpha_3}$ fully symmetric in $\alpha_1, \alpha_2, \alpha_3$.

Finally all is reduced to diagonalize the matrix $\mathcal{E}_{\alpha_1, \alpha_2, \alpha'_1, \alpha'_2}$ defined by

$$\Phi = \frac{3}{2} \sum_{\alpha_1, \alpha_2, \alpha'_1, \alpha'_2, \alpha_3} c_{\alpha_1, \alpha_2, \alpha_3}^* c_{\alpha'_1, \alpha'_2, \alpha_3} \mathcal{E}_{\alpha_1, \alpha_2, \alpha'_1, \alpha'_2}, \quad (13)$$

multiplied by the identity for the third gluon and symmetrized in $(1, 2, 3)$ and $(1', 2', 3)$.

Our concrete choice of trial basis are the harmonic oscillator eigenfunctions:

$$\begin{aligned} \psi_{\alpha}(\mathbf{r}) &= \psi_{k,l}(z) \exp il\phi, \quad z = \ln r^2, \quad \psi_{k,l}(-\infty) = 0, \\ \xi_k(z) &= c_k H_k(z) \exp(-z^2/2) = (\partial + (1/2)|l|) \psi_{k,l}(z), \end{aligned} \quad (14)$$

with $H_k(z)$ the Hermite polynomials.

4 Numerical results

From our numerical experience the best results are obtained with $|l| \leq l_{max}$, $k = 0, 1, \dots, (l_{max} + 1 = r)$. They are shown in Table 1.

These results are related to the corresponding intercepts as

$$1 + \omega_{BFKL}(0) = 1 - \frac{N_c \alpha_s}{\pi} \epsilon_2, \quad 1 + \omega_{odd}(0) = 1 - \frac{N_c \alpha_s}{\pi} \frac{3}{2} \epsilon_3. \quad (15)$$

The dimension of the matrix grows, for the three-gluon case, from 12 ($r = 2$) to 3368 ($r = 6$). For $r = 1$ the potential energy U_{ik} vanishes and one gets the kinetic energy per gluon, i.e. $\epsilon_2 = \epsilon_3$ in this case. It can be seen that the convergence of the method is quite slow.

In Table 2 we compare our results to other calculations. Our lower bound is compatible with our previous argument of $\epsilon_3 = 0$ but weaker than that of [10], which gives

Table 1: Lowest eigenvalues of (8),(9) for the two- (ϵ_2) and three-gluon (ϵ_3) bound states, using our trial functions labelled by r . $-2.773 \simeq -4 \ln 2$ is the exact value for $\pi \omega_{BFKL}(0)/(N_c \alpha_s)$.

r	ϵ_2	ϵ_3
1	0.968	0.968
2	0.022	0.605
3	-0.475	0.454
4	-0.743	0.379
5	-0.912	0.331
6	-1.032	0.298
∞	-2.773	

a supercritical odderon; this discrepancy may have a numerical origin. As to the results of [8], they are in principle exact but contain some semiclassical approximation whose reliability is difficult for us to quantify. So our conclusion is that the odderon intercept lies in the interval $0.91 \div 1.79$ (for $N_c = 3$ and $\alpha_s = 0.2$).

To quantify the existing uncertainty in more practical terms, let us consider the process $\gamma p \rightarrow \eta_c p$, i.e. diffractive photoproduction of η_c . The corresponding cross-section has been estimated [18] to be $\sigma(\gamma p \rightarrow \eta_c p) = D \times (47 \div 100)$ pb, $D = (\bar{x})^{-2\omega_{odd}(0)}$. From the previous considerations, D (for $N_c = 3$ and $\alpha_s = 0.2$) lies in the range $0.3 \div 55000$ for HERA ($\bar{x} \simeq 10^{-3}$). Clearly this reaction, in case it could be studied experimentally, is very sensitive to the value of the odderon intercept and offers a good opportunity to measure it.

In conclusion we have studied the odderon singularity as solution of the BKP equation for three colour-symmetric gluons and given an argument for the odderon intercept

Table 2: Comparison of our results to other calculations of the odderon intercept.

	$-(3/2)\epsilon_3$	$1 + \omega_{odd}(0)$ for $N_c = 3, \alpha_s = 0.2$
Our result	≥ -0.45	≥ 0.91
Ref. [10]	≥ 0.37	≥ 1.07
Ref. [8]	2.41	1.46
Upper bound from [9]	≤ 4.16	≤ 1.79

to be equal to unity. Besides a variational method to compute it has been presented and its results compared to other calculations. The odderon intercept lies in the interval $1 - (N_c\alpha_s/\pi)0.45 \leq j_{odd}(0) = 1 + \omega_{odd}(0) \leq 1 + (N_c\alpha_s/\pi)4.16$.

Acknowledgments

The authors thank J. Bartels, P. Gauron, G. P. Korchemsky, L. N. Lipatov, B. Nicolescu and S. Wallon for useful discussions and the organizers for such a nice meeting. N. A. also thanks Dirección General de Investigación Científica y Técnica of Spain for financial support.

References

- [1] P. D. B. Collins, *Regge Theory and High Energy Physics* (Cambridge University Press, Cambridge, 1977).
- [2] A. Donnachie and P.V. Landshoff, Nucl. Phys. **B267**, 690 (1986).
- [3] L. Lukaszuk and B. Nicolescu, Lett. Nuovo Cimento **8**, 405 (1973).
- [4] J.-R. Cudell, K. Kang and S. K. Kim, preprint SNUTP-95-103 (hep-ph/9601336); P. Gauron, B. Nicolescu and O. V. Selyugin, Phys. Lett. **B397**, 305 (1997).
- [5] E. A. Kuraev, L. N. Lipatov and V. S. Fadin, Sov. Phys. JETP **44**, 433 (1976); **45**, 199 (1977); Ya. Ya. Balitsky and L. N. Lipatov, Sov. J. Nucl. Phys. **28**, 822 (1978).
- [6] J. Bartels, Nucl. Phys. **B175**, 365 (1980); J. Kwieciński and M. Praszalowicz, Phys. Lett. **B94**, 413 (1980).
- [7] L. N. Lipatov, JETP Lett. **59**, 571 (1994); L. D. Faddeev and G. P. Korchemsky, Phys. Lett. **B342**, 311 (1995); Z. Maassarani and S. Wallon, J. Phys. **A28**, 6423 (1995).
- [8] G. P. Korchemsky, Nucl. Phys. **B443**, 255 (1995); **B462**, 333 (1996); in *Proceedings of the 28th International Conference on High Energy Physics*, eds. Z. Ajduk and A. K. Wroblewski (World Scientific, Singapore, 1997), p. 713.
- [9] P. Gauron, L. N. Lipatov and B. Nicolescu, Phys. Lett. **B260**, 407 (1991).
- [10] P. Gauron, L. N. Lipatov and B. Nicolescu, Z. Phys. **C63**, 253 (1994).
- [11] N. Armesto and M. A. Braun, Santiago preprint US-FT/9-94 (hep-ph/9410411); US-FT/7-96 (hep-ph/9603218) (Z. Phys. **C** to appear).

- [12] L. N. Lipatov in *Perturbative Quantum Chromodynamics*, ed. A. H. Mueller (World Scientific, Singapore, 1989), p. 411; J. Bartels, Phys. Lett. **B298**, 204 (1993); Z. Phys. **C60**, 471 (1993).
- [13] G. Camici in these proceedings; L. N. Lipatov in these proceedings.
- [14] A. H. Mueller, Phys. Lett. **B396**, 251 (1997); G. Camici and M. Ciafaloni, Phys. Lett. **B395**, 118 (1997).
- [15] T. Jaroszewicz, Acta Phys. Pol. **B11**, 965 (1980).
- [16] L. N. Lipatov, Sov. J. Nucl. Phys. **23**, 338 (1976).
- [17] L. N. Lipatov, Sov. Phys. JETP **63**, 904 (1986).
- [18] J. Czyżewski *et al.*, Phys. Lett. **B398**, 400 (1997); R. Engel *et al.*, preprint DESY 97-139 and NTZ 16/97 (hep-ph/9707362).